

LECTURE 7FOURIER SERIES

①

DEF Let $f, g: [-\pi, \pi] \rightarrow \mathbb{R}$ be CTS. (but not necessarily periodic)

Define the L^2 -INNER PRODUCT of f and g by

$$\langle f, g \rangle := \frac{1}{\pi} \int_{-\pi}^{\pi} f(x)g(x) dx$$

and the NORM of f by

$$\|f\| := \sqrt{\langle f, f \rangle}.$$

LEMMA The functions $1, \cos x, \sin x, \cos 2x, \sin 2x, \dots$ satisfy are PERIODIC on $[-\pi, \pi]$ and satisfy

$$\langle \cos kx, \cos lx \rangle = 0 = \langle \sin lx, \sin kx \rangle, \quad \begin{matrix} l \neq k \\ l, k \geq 0 \end{matrix}$$

$$\langle \cos kx, \sin lx \rangle = 0 \quad \forall l, k \geq 0$$

$$\|1\| = \sqrt{2}, \quad \|\cos kx\| = 1 = \|\sin kx\| \quad \forall k \geq 0.$$

PF

$$\textcircled{1} \quad \langle \cos kx, \sin lx \rangle = \int_{-\pi}^{\pi} (\cos kx) (\sin lx) dx = 0$$

$\begin{matrix} \uparrow & & \text{EVEN} \times \text{ODD} \\ \text{EVEN} & & \text{ODD} \end{matrix}$

(3)

$$\textcircled{2} \quad \frac{1}{\pi} \int_{-\pi}^{\pi} \cos(kx) \cos(lx) dx$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} [\cos((k+l)x) + \cos((k-l)x)] dx$$

$$= \begin{cases} \frac{1}{2\pi} \left[\frac{\sin((k+l)x)}{k+l} + \frac{\sin((k-l)x)}{k-l} \right]_{-\pi}^{\pi} = 0 & \text{if } k \neq l \\ \frac{1}{2\pi} \int_{-\pi}^{\pi} 1 dx = 1 & k = l. \end{cases}$$

□

SUPPOSE THAT

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

Then formally (without worrying whether series converge, etc):

$$\langle f(x), \cos(lx) \rangle = \frac{a_0}{2} \langle 1, \cos(lx) \rangle$$

$$+ \sum_{k=1}^{\infty} a_k \langle \cos(kx), \cos(lx) \rangle$$

$$+ b_k \langle \sin(kx), \cos(lx) \rangle$$

(3)

$$= \begin{cases} \frac{a_0}{2} & \text{if } l=0 \\ a_l & \text{if } l \neq 0 \end{cases}$$

Similarly $\langle f(x), \sin(lx) \rangle = b_l$.

DEF The FOURIER SERIES of $f: [-\pi, \pi] \rightarrow \mathbb{R}$

is

$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

where

$$a_k = \langle f, \cos kx \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(kx) dx \quad k=0, 1, 2, \dots$$

$$b_k = \langle f, \sin kx \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(kx) dx \quad k=1, 2, \dots$$

NOTE

- ① NO GUARANTEE that F.S. converges, or that if it does that ~~of~~ its value at x equals $f(x)$.

Hence notation $\sim$ instead of $=$.

- ② If the F.S. of f converges, it may NOT be C.B or differentiable (term by term) even if f is.

(4)


Ex 5

① [See over p76] $f(x) = x$.
 ✓

$$x \sim 2 \left(\sin x - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \frac{\sin 4x}{4} + \dots \right)$$

$$= 2 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin(kx)$$

② $f(x) = \text{sign}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$



$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} \text{sign}(x) dx = 0 \quad \text{by Symmetry.}$$

~~$$b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} \text{sign}(x) \sin(kx) dx$$~~

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} \text{sign}(x) \cos(kx) dx = 0$$

ODD $\times$ EVEN = ODD

BUT

$$b_k = \frac{1}{\pi} \int_{-\pi}^0 -\sin kx dx + \frac{1}{\pi} \int_0^{\pi} \sin kx dx$$

$$= \frac{2}{\pi} \int_0^{\pi} \sin(kx) dx = \frac{2}{\pi} \left[-\frac{\cos(kx)}{k} \right]_0^{\pi} \quad k=1, 2, \dots$$

(5)

$$b_k = \frac{2}{\pi} \left[\frac{1}{k} - \frac{\cos kx}{k} \right]$$

$$= \begin{cases} 0 & \text{if } k \text{ even} \\ \frac{4}{\pi k} & \text{if } k \text{ odd } (k=2n+1) \end{cases}$$

So

$$\text{sign}(x) \sim \sum_{n=0}^{\infty} \frac{4}{\pi(2n+1)} \sin[(2n+1)x]$$

DEF The n -th PARTIAL SUM of the FS of f
is

$$S_n(x) = \frac{a_0}{2} + \sum_{k=0}^n [a_k \cos kx + b_k \sin kx]$$

We say that the FS of f CONVERGES at x
if

$$\lim_{n \rightarrow \infty} S_n(x) = \hat{f}(x) \quad \text{I.}$$

WE WOULD LIKE $f(x) = \hat{f}(x) \quad \forall x \in [-\pi, \pi]$

But There is no guarantee This holds,

PERIODIC EXTENSIONS



FOURIER SERIES OF f

We start with $f: [-\pi, \pi] \rightarrow \mathbb{R}$.

We obtain

$$S_n(x) = \frac{a_0}{2} + \sum_{k=1}^n a_k \cos kx + b_k \sin kx$$

~~NOTICE~~ NOTICE

$S_n: \mathbb{R} \rightarrow \mathbb{R}$ is periodic:

$$S_n(x+2\pi) = S_n(x).$$

So if
have

$$\tilde{f}(x) = \lim_{n \rightarrow \infty} S_n(x) \quad \exists \text{ we must}$$

$$\tilde{f}(x+2\pi) = \tilde{f}(x), \quad \tilde{f}: \mathbb{R} \rightarrow \mathbb{R}$$

PERIODIC EXTENSION OF f

Given $f: [-\pi, \pi] \rightarrow \mathbb{R}$ we can define

$\tilde{f}_{\text{PER}}: \mathbb{R} \rightarrow \mathbb{R}$ periodic of period 2π so that


$$\tilde{f}_{\text{PER}}(x) = f(x) \quad \forall x \in (-\pi, \pi].$$

MAYBE


$\tilde{f} = \tilde{f}_{\text{PER}}$ for nice enough f ?

7

EX ① $f(x) = x$ on $[-\pi, \pi]$ has



② $f(x) = \text{sign}(x)$ on $[-\pi, \pi]$ has



§

DEF $f : [a, b] \rightarrow \mathbb{R}$ is PIECEWISE CONTINUOUS (PWCS) if

① f is defined and continuous on $[a, b]$ except possibly at a finite # of points


$$a \leq x_1 < x_2 < \dots < x_n \leq b,$$

② At each discontinuity x_k we require

$$\lim_{x \rightarrow x_k^-} f(x) =: f(x_k^-) \exists$$

$$\lim_{x \rightarrow x_k^+} f(x) =: f(x_k^+) \exists.$$

EX



DEF $f : [a, b] \rightarrow \mathbb{R}$ is PIECEWISE C^1 if

① f is defined, CTS, C^1 (continuously differentiable) on $[a, b]$ except possibly at finite # pts $a \leq x_1 < x_2 < \dots < x_n \leq b$


⑨

⑥ At each x_k we require

$$\lim_{x \rightarrow x_k^\pm} f(x) =: f(x_k^\pm) \exists$$


$$\lim_{x \rightarrow x_k^\pm} f'(x) =: f'(x_k^\pm) \exists.$$

EX



NOTE ① $f(x) = \sin\left(\frac{1}{x}\right)$ on $[-1, 1]$ is not pbc

as $\lim_{x \rightarrow 0^\pm} \sin\left(\frac{1}{x}\right)$ DNE



$$\textcircled{2} \quad f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

is CTS on $[-1, 1]$ but not PW C^1

as

$$f'(x) = \sin\left(\frac{1}{x}\right) - \frac{1}{x} \cos x, \quad x \neq 0$$

has $\lim_{x \rightarrow 0^\pm} f'(x)$ DNE.

