

LECTURE 8

FOURIER SERIES, PART II


①

THM ON POINTWISE CONVERGENCE OF F.S

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a 2π -periodic, piecewise C^1 function.

Then at any $x \in \mathbb{R}$, the Fourier Series of f converges to

$$\begin{cases} f(x) & \text{if } f \text{ is CT at } x \\ \frac{1}{2} [f(x^+) + f(x^-)] & \text{if } f \text{ has JUMP DISCTY at } x \end{cases}$$




EX LET $f: [-\pi, \pi] \rightarrow \mathbb{R}$ be defined by

$$f(x) = x \quad \text{for } -\pi \leq x \leq \pi$$

Extend f to a 2π -periodic PW C^1 f^n ,
which we will ~~stop~~ call f_{PER}

(2)



$$\tilde{f}_{\text{PER}}(x) = x - 2m\pi \quad \text{for } (2m-1)\pi \leq x < (2m+1)\pi$$

where $m \in \mathbb{Z}$.

CHECK

- Slope = 1
- Crosses x-axis at $x = 2m\pi$.

LAST TIME: Fourier Series of f is


$$x \sim 2 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin(kx)$$

SO PW CONVERGE THUS SATIS

$$2 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin(kx) = \begin{cases} x - 2m\pi & \text{IF } (2m-1)\pi < x < (2m+1)\pi \\ & \text{FOR SOME } m \\ 0 & \text{IF } x = (2n-1)\pi \\ & \text{FOR SOME } n \end{cases}$$

$$=: \tilde{f}(x)$$


(3)

NOTICE

① $\tilde{f}(0) = \infty$ ✓ $\tilde{f}(\pi) = 0$.

② When $x = \pi/2$ we have

$$\frac{\pi}{2} = 2 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin\left(\frac{k\pi}{2}\right)$$



S_0

$$\frac{\pi}{4} = \sum_{m=0}^{\infty} \frac{(-1)^{2m+2}}{2m+1} (-1)^m = \sum_{m=0}^{\infty} \frac{(-1)^m}{2m+1}$$

$$= 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

Although the FS converges at each x , The convergence is not UNIFORM.

We have $\tilde{f}(x) = \lim_{n \rightarrow \infty} S_n(x)$ where

$S_n = n\pi$ partial sum of FS.

(4)

But the # of terms, n , need to ensure error is less than ϵ ,

$$|\hat{f}(x) - s_n(x)| < \epsilon$$

depends on x :

Converge is only PW

$$\forall x \forall \epsilon > 0 \exists N = N(\epsilon, x) : \forall n \geq N |s_n(x) - \hat{f}(x)| < \epsilon$$






Rather than UNIFORM

$$\forall \epsilon > 0 \exists N = N(\epsilon) : \forall x, \forall n \geq N |s_n(x) - \hat{f}(x)| < \epsilon$$

GIBB'S PHENOMENON

- Away from jump discontinuities, series converges.
- Near jumps $\exists$ overshoot of $\sim 9\%$ of jump magnitude
- As $n \uparrow$ width of overshoot region $\downarrow$, but magnitude persists, indept of n .

Fourier Series of Sawtooth Function and Gibbs Phenomenon



(6)

FOURIER COSINE SERIES

Let $f: [-\pi, \pi] \rightarrow \mathbb{R}$ be even.

Then
$$b_k = \int_{-\pi}^{\pi} f(x) \sin(kx) dx = 0 \quad \forall k$$

EVEN $\times$ ODD

So
$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx)$$
 FOURIER
COSINE SERIES

WITH
$$a_k = \frac{2}{\pi} \int_0^{\pi} f(x) \cos(kx) dx$$

FOURIER SINE SERIES

Let $f: [-\pi, \pi] \rightarrow \mathbb{R}$ be ODD

Then
$$a_k = \int_{-\pi}^{\pi} f(x) \cos(kx) dx = 0 \quad \forall k$$

ODD $\times$ EVEN

So
$$f(x) \sim \sum_{k=1}^{\infty} b_k \sin(kx)$$
 FOURIER
SINE SERIES

WITH
$$b_k = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(kx) dx$$

CONSTRUCTION

- ① Let $f: [0, \pi] \rightarrow \mathbb{R}$. Then f has a unique
 EVEN extension to $\tilde{f}_{\text{EVEN}}: [-\pi, \pi] \rightarrow \mathbb{R}$
 defined by setting

$$\tilde{f}_{\text{EVEN}}(x) = f(x) \quad \text{for } 0 \leq x \leq \pi$$

$$\tilde{f}_{\text{EVEN}}(x) = f(-x) \quad \text{for } -\pi \leq x < 0.$$

(7)

We can further extend $\tilde{f}_{\text{EVEN}}$ to a 2π -PERIODIC
EVEN function $\tilde{f}_{\text{EVEN, PER}} : \mathbb{R} \rightarrow \mathbb{R}$.

Fact


$$\tilde{f}_{\text{EVEN, PER}}(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx)$$

with

$$a_k = \frac{2}{\pi} \int_0^{\pi} f(x) \cos(kx) dx$$

The Fourier Cosine Series of f represents the even 2π -periodic extension of f .

EX



So $\tilde{f}_{\text{EVEN, PER}}(x) = 1 \quad \forall x \in \mathbb{R}$.

We have

$$a_k = \frac{2}{\pi} \int_0^{\pi} 1 \cos(kx) dx = \begin{cases} 2 & k=0 \\ 0 & k>0 \end{cases}$$

So Fourier Cosine Series of $f = \frac{a_0}{2} = 1$ ✓

8

(2) Let $f: [0, \pi] \rightarrow \mathbb{R}$, ^{WITH $f(0)=0$} Then f has a unique odd extension

$$\tilde{f}_{\text{ODD}}: [-\pi, \pi] \rightarrow \mathbb{R}$$

defined by

$$\tilde{f}_{\text{ODD}}(x) = f(x), \quad 0 \leq x \leq \pi$$

$$\tilde{f}_{\text{ODD}}(x) = -f(-x), \quad -\pi \leq x < 0.$$

We can further extend $\tilde{f}_{\text{ODD}}$ to a 2π -periodic odd f^n

$$\tilde{f}_{\text{ODD, PER}}: \mathbb{R} \rightarrow \mathbb{R}.$$




FACT

$$\tilde{f}_{\text{ODD, PER}}(x) \sim \sum_{k=1}^{\infty} b_k \sin(kx)$$

$$b_k = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(kx) dx$$

The Fourier sine series represents the odd 2π -periodic extension of f .

(9)



$$b_k = \frac{2}{\pi} \int_0^{\pi} 1 \sin(kx) dx$$

$$= \frac{2}{\pi} \left[\frac{-1}{k} \cos(kx) \right]_0^{\pi} = \frac{-2}{\pi k} [\cos(k\pi) - 1]$$

$$b_k = \begin{cases} 0 & \text{if } k = 2n, \text{ EVEN} \\ \frac{4}{\pi(2n+1)} & \text{if } k = 2n+1, \text{ ODD} \end{cases}$$

$$S_0 \quad \tilde{f}_{\text{ODD,PER}}(x) \sim \sum_{n=0}^{\infty} \frac{4}{\pi(2n+1)} \sin[(2n+1)x]$$

DIFFERENTIATION OF FOURIER SERIES

Suppose

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(kx) + b_k \sin(kx).$$

We would like to have

$$f'(x) = \sum_{k=1}^{\infty} -ka_k \sin(kx) + kb_k \cos(kx).$$

wherever f is differentiable.

"DIFFERENTIATE TERM BY TERM"

TROUBLING EXWe just proved that for $0 < x < \pi$ we have

$$1 = \sum_{n=0}^{\infty} \frac{4}{\pi(2n+1)} \sin[(2n+1)x]$$

BUT

$$0 = \frac{d}{dx}(1) \neq \sum_{n=0}^{\infty} \frac{4}{\pi} \cos[(2n+1)x]$$

↑
DIVERGES for $0 < x < \pi$ So we need to impose some extra conditions on f to allow us to differentiate F.S. term by term.IDEA To ensure PS of f' CONVS we need f' PW C^1 .

THM ON DIFFERENTIATION OF F.S.

Suppose that $f: [-\pi, \pi]$ has a CTS, 2π -PERIODIC piecewise C^2 extension to $\mathbb{R}$.

$$\text{If } f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

$$\text{Then } f'(x) \sim \sum_{k=1}^{\infty} k b_k \cos kx - k a_k \sin kx$$

EXS


① Our previous example was not CTS so Thm does not apply.

② Let $f(x) = |x|$.

on $[-\pi, \pi]$:

$$f'(x) = \text{sign}(x)$$

$$= \begin{cases} +1 & x > 0 \\ -1 & x < 0 \end{cases}$$



f has CTS 2π -periodic, PWC 2 ext n to $\mathbb{R}$ ✓

UB ✓

$$|x| \sim \frac{\pi}{2} - \frac{4}{\pi} \left[\cos x + \frac{\cos 3x}{9} + \frac{\cos 5x}{25} + \dots \right]$$

$$\text{And so } \text{sign}(x) \sim \frac{4}{\pi} \left[\sin x + \frac{\sin 3x}{3} + \frac{\sin 5x}{5} + \dots \right]$$

which UB ✓ is FS of $\text{sign}(x)$.

CHANGE OF SCALE

12

Let $f: [-L, L] \rightarrow \mathbb{R}$.

Define $s: [-\pi, \pi] \rightarrow [-L, L]$ by

$$s(y) = \frac{L}{\pi} y$$

and set

$$F(y) = f(s(y)) = f\left(\frac{L}{\pi} y\right) = f(x) \quad x = \frac{L}{\pi} y$$

So that

$$F: [-\pi, \pi] \rightarrow \mathbb{R}.$$

We have

$$F(y) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos(ky) + b_k \sin(ky)$$

with

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} F(y) \cos(ky) dy$$

$$b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} F(y) \sin(ky) dy.$$

So

$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos\left(\frac{k\pi x}{L}\right) + b_k \sin\left(\frac{k\pi x}{L}\right)$$

with

$$a_k = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{k\pi x}{L}\right) dx \quad dy = \frac{\pi}{L} dx$$

$$b_k = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{k\pi x}{L}\right) dx$$