

COMPLEX FOURIER SERIESEVLER'S FORMULA

$$e^{ikx} = \cos(kx) + i \sin(kx)$$

$$e^{-ikx} = \cos(kx) - i \sin(kx)$$

So

$$\cos(kx) = \frac{1}{2} (e^{ikx} + e^{-ikx})$$

$$\sin(kx) = \frac{1}{2i} (e^{ikx} - e^{-ikx})$$

LET'S PLAY IDEA CALCULATIONS OFTEN EASIER WITH
COMPLEX EXPS THAN WITH TRIG FNS

$$a_k \cos kx + b_k \sin kx$$

$$= \frac{1}{2} a_k (e^{ikx} + e^{-ikx}) + \frac{1}{2i} b_k (e^{ikx} - e^{-ikx})$$

$$= \frac{1}{2} (a_k - i b_k) e^{ikx} + \frac{1}{2} (a_k + i b_k) e^{-ikx}$$

So set

$$c_k = \frac{1}{2} (a_k - i b_k) \quad k > 0$$

$$c_{-k} = \frac{1}{2} (a_k + i b_k) \quad k > 0$$

$$c_0 = \frac{1}{2} a_0$$

(2)

Then

$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

$$= c_0 e^{i0x} + \sum_{k=1}^{\infty} c_k e^{ikx} + c_{-k} e^{-ikx}$$

$$= \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

and

$$c_k = \frac{1}{2} (a_k - ib_k)$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) [\cos kx - i \sin kx] dx$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \overline{e^{ikx}} dx$$

$$c_k = \langle f, e^{ikx} \rangle$$

SUMMARY

① Let $f, g: [-\pi, \pi] \rightarrow \mathbb{C}$. Define HERMITIAN L^2 -INNER PRODUCT by

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \overline{g(x)} dx$$

(3)

③ Norm of f : $\|f\| = \sqrt{\langle f, f \rangle} = \sqrt{\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x)|^2 dx}$.

③ $\{e^{ikx} / k \in \mathbb{Z}\}$ is ONB in that

$$\langle e^{ikx}, e^{ilx} \rangle = \delta_{kl}.$$

PF

③ $\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikx} \overline{e^{ilx}} dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(k-l)x} dx$

$$= \begin{cases} \frac{1}{2\pi} \left[\frac{e^{i(k-l)x}}{i(k-l)} \right]_{-\pi}^{\pi} = 0 & \text{if } k \neq l \\ \frac{1}{2\pi} \int_{-\pi}^{\pi} 1 dx = 1 & \text{if } k = l. \end{cases}$$

④ Let $f: [-\pi, \pi] \rightarrow \mathbb{C}$. The complex FS of f is

$$f(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

where

$$c_k = \langle f, e^{ikx} \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx$$

(4)

EX $f(x) = e^{ax}$, $a \neq 0$.

$$c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ax} e^{-ikx} dx$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{(a-ik)x} dx$$

$$= \frac{1}{2\pi} \left[\frac{e^{(a-ik)x}}{a-ik} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{2} \frac{e^{(a-ik)\pi} - e^{-(a-ik)\pi}}{(a-ik)\pi} \quad e^{ik\pi} = (-1)^k$$

$$= (-1)^k \frac{e^{a\pi} - e^{-a\pi}}{2\pi (a-ik)}$$

$$= (-1)^k \frac{\sinh(a\pi)}{\pi (a-ik)} \quad \sinh x = \frac{e^x - e^{-x}}{2}$$

$$= (-1)^k \frac{(a+ik) \sinh(a\pi)}{\pi (a^2+k^2)}$$

So $e^{ax} \sim \frac{\sinh(a\pi)}{\pi} \sum_{k=-\infty}^{\infty} \frac{(-1)^k (a+ik)}{a^2+k^2} e^{ikx} \quad (*)$

(5)

NOTICE If set $a=1$, $x=\infty$ get

$$\frac{\pi}{\sinh(\pi)} = \sum_{k=-\infty}^{\infty} \frac{(-1)^k (1+ik)}{1+k^2}$$

$$= \sum_{k=-\infty}^{\infty} \frac{(-1)^k}{1+k^2} + i \sum_{k=-\infty}^{\infty} \frac{(-1)^k k}{1+k^2}$$

$$= 1 + 2 \sum_{k=1}^{\infty} \frac{(-1)^k}{1+k^2}$$

$$\text{as } \sum_{k=-\infty}^{-1} \frac{(-1)^k k}{1+k^2} = - \sum_{l=1}^{\infty} \frac{(-1)^l l}{1+l^2}$$

$(l=-k)$

$$S_0 \quad \sum_{k=-\infty}^{\infty} \frac{(-1)^k k}{1+k^2} = 0 \quad \text{HOLDS!}$$

EXERCISE

Convert $(*)$ to F.S. involving TRG p_n .

(6)

POINTWISE + UNIFORM CONVERGENCE

FOR PROOFS SEE AN ANALYSIS COURSE LIKE MATH 4301/4302

DEF ① Let $f_n : [a, b] \rightarrow \mathbb{R}$ be a sequence of f^n .
 We say $f_n \rightarrow f$ POINTWISE if

$$\forall x \in [a, b] \quad f_n(x) \rightarrow f(x) \text{ as } n \rightarrow \infty$$

is if $\forall x \in [a, b], \quad \forall \varepsilon > 0, \exists N : \forall n \geq N$

$$|f_n(x) - f(x)| < \varepsilon.$$

$$N = N(\varepsilon, x)$$

② Let $f_n : [a, b] \rightarrow \mathbb{R}$. We say $f_n \rightarrow f$ UNIFORMLY ON $[a, b]$ if

$$\forall \varepsilon > 0, \exists N : \forall x \in [a, b], \forall n \geq N$$

$$|f_n(x) - f(x)| < \varepsilon$$

$$N = N(\varepsilon)$$

EX

$$f_n(x) = x^n \quad \text{on } [0, 1]$$

$$f(x) = \begin{cases} 0 & \text{if } x < 1 \\ 1 & \text{if } x = 1 \end{cases}$$

Then

$f_n \rightarrow f$ POINTWISE but NOT UNIFORMLY.

(7)

as for $x < 1$

$$|f_n(x) - f(x)| = |x^n - 0| = |x|^n < \varepsilon$$

$$\Leftrightarrow n \log |x| < \log \varepsilon \quad (\log \text{ is } \uparrow)$$

$$\Leftrightarrow n > \frac{\log \varepsilon}{\log |x|} \quad \text{as } \log |x| < 0 \text{ for } 0 < x < 1$$

$$\text{Set } N(\varepsilon, x) = \frac{\log \varepsilon}{\log |x|}$$

$$\text{As } x \rightarrow 1, \quad N(\varepsilon, x) \rightarrow \infty.$$

So don't expect uniform converge.

THM A If $f_n \rightarrow f$ UNIF on $[a, b]$

and each f_n is CT on $[a, b]$ Then

f is CT on $[a, b]$

NOTE EX above shows PW converge not enough to ensure limit is CT.

⑧

DEF Let $u_k : [a, b] \rightarrow \mathbb{R}$, $k = 1, 2, \dots$

and
$$S_n(x) = \sum_{k=1}^n u_k(x).$$

We say $\sum_{k=1}^{\infty} u_k$ CONVERGES (PW/UNIF) if

$$S_n \rightarrow S \quad \text{PW/UNIF}$$

WEIERSTRASS M-TEST

Suppose $|u_k(x)| \leq m_k \quad \forall x \in [a, b]$
[Each u_k is bounded]

If $\sum_{k=1}^{\infty} m_k$ CONVS.

Then $\sum_{k=1}^{\infty} u_k(x) = f(x)$

CONVS UNIFLY + ABSOLUTELY.

Consequently (by Thm A) if each u_k is CT, so is f .

⑨

We can manipulate UNIFLY CONVFT series just like finite sums.

THM B

Suppose

$$f(x) = \sum_{k=1}^{\infty} u_k(x) \quad \text{CONVFS UNIF}$$

Then

$$\textcircled{1} \quad \int g(x) f(x) = \sum_{k=1}^{\infty} g(x) u_k(x)$$

for any bounded $f^n g$.

$$\textcircled{2} \quad \int_a^b f(y) dy = \int_a^b \left[\sum_{k=1}^{\infty} u_k(y) \right] dy$$

$$= \sum_{k=1}^{\infty} \int_a^b u_k(y) dy$$

- Can integrate series term by term

THM C

Suppose

$$f(x) = \sum_{k=1}^{\infty} u_k(x) \quad \text{POINTWISE}$$

and

$$\sum_{k=1}^{\infty} u'_k(x) =: g(x) \quad \text{CONVS UNIF}$$

Then

$$\sum_{k=1}^{\infty} u_k(x) \quad \text{CONVS UNIF} \quad \text{and}$$

(10)

$$f'(x) = g(x)$$

$$\frac{d}{dx} \left[\sum_{k=-\infty}^{\infty} u_k(x) \right] = \sum_{k=-\infty}^{\infty} \frac{du_k}{dx}(x)$$

- TERM BY TERM DIFFN.

Thm D

Suppose $f(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$

IF $\sum_{k=-\infty}^{\infty} |c_k|^2 < \infty$

Then the F.S. CONVS UNIF to a CTS $\tilde{f}$

WITH same FS coefficients as f :

$$c_k = \langle \tilde{f}, e^{ikx} \rangle = \langle f, e^{ikx} \rangle$$

IF

$$|c_k e^{ikx}| \leq c_k \quad \text{as } |e^{ikx}| = 1$$

So $\tilde{f}(x) = \sum c_k e^{ikx}$ CONVS by Weierstrass Test.

(11)

Then

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \hat{f}(x) e^{-ikx} dx$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left[\sum_k c_k e^{ikx} \right] e^{-ikx} dx$$

$$\text{THM B1} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left[\sum_k c_k e^{i(k-l)x} \right] dx$$

$$\text{THM B2} = \frac{1}{2\pi} \sum_k c_k \int_{-\pi}^{\pi} e^{i(k-l)x} dx$$

$$= \sum_k c_k \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(k-l)x} dx$$

$$= \sum_k c_k \delta_{kl} = c_l$$

□


THM E

Let f be 2π -periodic and piecewise C^1 .

If f is CTB on subinterval (a, b)

Then F.S. of f convs unif to f

on any closed subinterval $[a+\delta, b-\delta]$ of (a, b) .

EX

F.S. converges UNIF to $f(x) = x$ on $[-\pi+\delta, \pi-\delta]$

- Related to Gibbs Phenomenon ~~is~~ only a problem at discontinuities $x = \pm\pi$.

— 0 —

THM F

Suppose

$$f(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

Then

①

$$\sum_{k=-\infty}^{\infty} |c_k|^2 \leq \|f\|^2$$

BESELS

INEQUALITY

②

Hence $c_k \rightarrow 0$ as $k \rightarrow \pm\infty$.

Also $a_k \rightarrow 0, b_k \rightarrow 0$ as $k \rightarrow \infty$
 as $a_k = c_k + c_{-k}, b_k = i(c_k - c_{-k})$

PF Let $\psi_k(x) = e^{ikx}$ and

$$S_n = \sum_{|k| \leq n} c_k \psi_k$$

NOW

$$\textcircled{A} \quad \|S_n\|^2 = \langle S_n, S_n \rangle$$

$$= \left\langle \sum_{|k| \leq n} c_k \psi_k, \sum_{|l| \leq n} c_l \psi_l \right\rangle$$

$$= \sum_{|k|, |l| \leq n} c_k \bar{c}_l \langle \psi_k, \psi_l \rangle$$

$$= \sum_{|k|, |l| \leq n} c_k \bar{c}_l \delta_{kl} = \sum_{|k| \leq n} |c_k|^2$$

$$\textcircled{B} \quad \langle f, S_n \rangle = \left\langle f, \sum_{|k| \leq n} c_k \psi_k \right\rangle$$

$$= \sum_{|k| \leq n} \bar{c}_k \langle f, \psi_k \rangle$$

$$= \sum_{|k| \leq n} |c_k|^2 = \|S_n\|^2$$

$$\textcircled{c} \quad 0 \leq \|f - s_n\|^2$$

$$= \langle f - s_n, f - s_n \rangle$$

$$= \|f\|^2 + \|s_n\|^2 - 2\langle f, s_n \rangle$$

$$\textcircled{1}, \textcircled{2} \quad \|f\|^2 = \sum_{|k| \leq n} |c_k|^2 \quad \checkmark$$

$$\text{So} \quad \sum_{|k| \leq n} |c_k|^2 \leq \|f\|^2$$

$$\textcircled{d} \quad a_n = \sum_{|k| \leq n} |c_k|^2 \quad \text{is } \uparrow \text{ seq, bounded}$$

$$\text{above.} \quad \text{So} \quad a_n \rightarrow a_\infty = \sum_{k=-\infty}^{\infty} |c_k|^2 \quad \text{convs.}$$

$$\text{So} \quad c_k \rightarrow 0 \quad \text{must hold as } k \rightarrow \pm\infty$$

□