

Math 2415

Problem Section #1

Make sure you do some problems from each section.

12.1: 3D Coordinate Systems

1. Draw a rectangular box with the origin and the point $(1, 2, 3)$ as opposite vertices and faces parallel to the coordinate planes. Label each vertex with its coordinates. Find the length of the diagonal of the box.
2. (a) What does the equation $x = 2$ represent in \mathbb{R}^2 ? Sketch!
(b) What does the equation $x = 2$ represent in \mathbb{R}^3 ? Sketch!
(c) What does the equation $z = 1$ represent in \mathbb{R}^3 ? Sketch!
(d) Describe the set of all points, (x, y, z) , in \mathbb{R}^3 for which $x = 2$ and $z = 1$. Sketch!
3. For what values of b and c do the points $(1, 2, 3)$, $(4, 5, 1)$, and $(10, b, c)$ all lie on the same line?
4. (a) Find the equation of the sphere with center $(1, 3, 5)$ and radius 4.
(b) What is the intersection of this sphere with the xz -plane? Argue algebraically and geometrically.
(c) What would the radius of the sphere have to be for the intersection of the sphere and the xz -plane to be a single point. What are the coordinates of this point?


12.2: Vectors

1. Do not use coordinate representations of vectors to solve this problem. Just draw pictures.
 - (a) Draw two vectors that are not parallel and label them \mathbf{a} and \mathbf{b} .
 - (b) Sketch the vector $\mathbf{a} + \mathbf{b}$
 - (c) Sketch the vector $\mathbf{a} - \frac{1}{2}\mathbf{b}$
 - (d) Sketch the vector $\frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$
2. Sketch a parallelogram and label the vertices A , B , C , and D going around counter-clockwise from the bottom left vertex. Let E be the point obtained by intersecting the two diagonals of the parallelogram. Make sure the side lengths of your parallelogram are not all equal, ie you did not draw a rhombus. The notation \overrightarrow{AB} refers to the displacement vector from the point A to the point B .
 - (a) Name all pairs of equal vectors in your sketch.
 - (b) Write each combination of vectors as a single vector: $\overrightarrow{AB} + \overrightarrow{BC}$, $\overrightarrow{AE} - \overrightarrow{EB}$, $2\overrightarrow{AB} + \overrightarrow{BD}$.
3. Find a unit vector parallel to $(1, 2, 3)$.
4. Let $\mathbf{a} = 3\mathbf{j} - 4\mathbf{k}$ and $\mathbf{b} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. Find

- (a) $\mathbf{a} + 2\mathbf{b}$
 - (b) $|\mathbf{b}|$
 - (c) $|\mathbf{a} - \mathbf{b}|$.
5. Suppose that $\mathbf{v} \in \mathbb{R}^2$ lies in the 2nd quadrant, makes an angle of 120° with the positive x -axis, and has length $|\mathbf{v}| = 2$. Find the coordinates of \mathbf{v} .


12.3: The Dot Product

1. Find $\mathbf{a} \cdot \mathbf{b}$ if
 - (a) $\mathbf{a} = (1, 2)$ and $\mathbf{b} = (-2, 3)$,
 - (b) $\mathbf{a} = 2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}$ and $\mathbf{b} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$,
 - (c) $|\mathbf{a}| = 3$, $|\mathbf{b}| = 4$, and the angle between \mathbf{a} and \mathbf{b} is 120° .
2. (a) Let $\mathbf{u} = (3, -2, 1)$ and $\mathbf{v} = (2, 4, -1)$.
 - (b) Find the angle between \mathbf{u} and \mathbf{v} to the nearest degree (use a calculator!)
 - (c) Find three nonzero vectors that are orthogonal to \mathbf{u} .
3. Find $\mathbf{u} \cdot \mathbf{v}$



4. Let C be the point on the line segment AB that is twice as far from A and it is from B , and let O denote the origin. Let $\mathbf{a} = \overrightarrow{OA}$, $\mathbf{b} = \overrightarrow{OB}$, and $\mathbf{c} = \overrightarrow{OC}$. [Recall that \overrightarrow{AB} refers to the displacement vector from the point A to the point B .]
 - (a) Make a sketch showing the relationships between all these points and vectors. Your sketch will help you solve the other parts of the problem.
 - (b) Express \overrightarrow{AB} in terms of \mathbf{a} and \mathbf{b} .
 - (c) Hence express \overrightarrow{AC} in terms of \mathbf{a} and \mathbf{b} .
 - (d) Hence find a formula for \mathbf{c} in terms of \mathbf{a} and \mathbf{b} .
5. What restrictions must be made on b so that the vector $2\mathbf{i} + b\mathbf{j}$ is orthogonal to (a) $-3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ and (b) \mathbf{k} ?
6. The cube in the figure has side length one with one vertex at the origin, O .
 - (a) Let Q be the vertex opposite O and let R be the midpoint of a face of the cube as shown in the figure. Find a formula for the vectors \overrightarrow{OQ} and \overrightarrow{OR} in terms of the vectors \mathbf{i} , \mathbf{j} , and \mathbf{k} .

- (b) Find the angle between the vectors \mathbf{i} and \overrightarrow{OQ} .
- (c) Find the angle between the vectors \overrightarrow{OQ} and \overrightarrow{OR} .



Extra Challenge Questions:

- Find the angle between a diagonal of a cube and a diagonal of one of its faces.
- We learned in class that $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos(\theta)$. Use this property to prove the Cauchy-Schwarz inequality

$$|\mathbf{a} \cdot \mathbf{b}| \leq |\mathbf{a}||\mathbf{b}|. \quad (1)$$

- The triangle inequality states that

$$|\mathbf{a} + \mathbf{b}| \leq |\mathbf{a}| + |\mathbf{b}|. \quad (2)$$

- Give a geometric interpretation of the triangle inequality.
- Use the fact that $|\mathbf{a} + \mathbf{b}|^2 = (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b})$ and the distribute law for the dot product to prove (2).

- The parallelogram law states that

$$|\mathbf{a} + \mathbf{b}|^2 + |\mathbf{a} - \mathbf{b}|^2 = 2|\mathbf{a}|^2 + 2|\mathbf{b}|^2. \quad (3)$$

- Give a geometric interpretation of the parallelogram law.
- Prove the parallelogram law.