

LAST NAME:	FIRST NAME:	CIRCLE:	Khoury 5:30pm	Coskunuzer 8:30am
CAYLEY	ARTHUR	Coskunuzer 11:30am	Zweck 1pm	Zweck 4pm

LOOK THEM UP ON WIKIPEDIA

1 /10	2 /10	3 /10	4 /10	5 /10
-------	-------	-------	-------	-------

6 /10	7 /10	8 /10	9 /10	10 /10	T /100
-------	-------	-------	-------	--------	--------

MATH 2415 Final Exam, Fall 2023

No books or notes! NO CALCULATORS! Show all work and give complete explanations. This 2 hours 45 mins exam is worth 100 points.

- (1) [10 pts] Find an equation of the form $Ax + By + Cz = D$ for the plane containing the points $(0, 1, 1)$, $(1, 0, 1)$, and $(1, 1, 0)$. Which of the following points lies on this plane: $P = (2, -1, 3)$, $Q = (3, -4, 3)$.

B C

$$\vec{u} = \vec{AB} = B - A = (1, -1, 0)$$

$$\vec{v} = \vec{AC} = C - A = (1, 0, -1)$$

$$\vec{n} = \vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{vmatrix}$$


$$= (1, 1, 1)$$

$$\vec{x}_0 = A = (0, 1, 1)$$

$$\text{So } (\vec{x} - \vec{x}_0) \cdot \vec{n} = 0 \text{ gives}$$

$$(x, y-1, z-1) \cdot (1, 1, 1) = 0$$

$$\boxed{x+y+z=2}$$



P:

$$2 - 1 + 3 = 4 \neq 2$$

P DOES NOT LIE ON PLANE

$$Q: 3 - 4 + 3 = 2$$

Q DOES LIE ON PLANE

(2) [10 pts]

(a) Find the volume of the parallelepiped determined by the vectors $\mathbf{u} = (1, 4, 2)$, $\mathbf{v} = (-1, 1, 4)$ and $\mathbf{w} = (5, 1, 2)$.



$$\text{VOL} = |(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \begin{vmatrix} 1 & 4 & 2 \\ -1 & 1 & 4 \\ 5 & 1 & 2 \end{vmatrix}$$

$$= |-2 - 4(-22) + 2(-6)| = \cancel{-102}$$

$$= \cancel{102}$$

†

(b) Make a sketch that shows how to project the vector $\mathbf{v} = \mathbf{i}$ onto the vector $\mathbf{w} = \mathbf{i} + \sqrt{3}\mathbf{j}$. Use your sketch to find the component of \mathbf{v} in the direction \mathbf{w} .



$\text{COMP}_{\mathbf{w}}(\mathbf{v}) = +|\overrightarrow{AP}|$ as $\text{PROJ}_{\mathbf{w}}(\mathbf{v})$ points in same dirn as \mathbf{w}


$= \frac{1}{2}$ from triangle II

(3) [10 pts]

(a) Sketch the surface $x^2 + y^2 - z^2 = 2$ for $0 \leq z \leq \sqrt{2}$.


This is a SURFACE OF REVOLUTION, in cyl coords:

$$r^2 - z^2 = 2$$



$$z=0: r=\sqrt{2}$$

$$\text{ASYMPTOTES: } z = \pm r$$



T

(b) Show that the line through the point $(1, 1, 0)$ in the direction of the vector $(-1, 1, \sqrt{2})$ lies on the surface in (a).

Parametrize the line:

$$\vec{r}(t) = \vec{p} + t\vec{v} = (1, 1, 0) + t(-1, 1, \sqrt{2})$$

$$(x, y, z) = \vec{r}(t) = (1-t, 1+t, \sqrt{2}t)$$

To show line lies on surface plug

$$x = 1-t, y = 1+t, z = \sqrt{2}t \text{ into } x^2 + y^2 - z^2 = 2.$$

$$\text{Well } x^2 + y^2 - z^2$$

$$= (1-t)^2 + (1+t)^2 - (\sqrt{2}t)^2$$

$$= 1 - 2t + t^2 + 1 + 2t + t^2 - 2t^2$$

$$= 2$$

(4) [10 pts] Find and classify all critical points of the function

$$f(x, y) = 3x^2y - 3y + y^3$$

This is a local optimization problem.

First find critical points of f :

$$\frac{\partial f}{\partial x} = 6xy \quad (1)$$

$$\frac{\partial f}{\partial y} = 3x^2 - 3 + 3y^2 \quad (2)$$

By (1) $x=0$ or $y=0$

CASE $x=0$

$$\text{By (2)} \quad y^2 = 1 \\ y = \pm 1$$

$$(x, y) = (0, \pm 1)$$

CASE $y=0$

$$\text{By (2)} \quad x^2 = 1$$


$$\text{So } x = \pm 1$$

$$(x, y) = (\cancel{0}, \pm 1) \quad (\pm 1, 0)$$

2nd DER TEST

$$D = \det \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix} = \det \begin{bmatrix} 6y & 6x \\ 6x & 6y \end{bmatrix} = 36(y^2 - x^2)$$

TEST	(x, y)	D	f_{xx}	CLASSIFICATION
(0, 0)	(0, 0)	0		
$(0, 1)$	$36 > 0$	$6 > 0$		Local M.W.
$(0, -1)$	$36 > 0$	$-6 < 0$		Local M.A.
$(1, 0)$	$-36 < 0$			SADDLE PT
$(-1, 0)$	$-36 < 0$			SADDLE PT




(5) [10 pts] Evaluate the double integral $\int_{x=0}^{x=1} \int_{y=\sqrt{x}}^{y=1} \sqrt{y^3 + 1} dy dx$. I

Since I have no idea how to do $\int \sqrt{y^3 + 1} dy$
lets switch order of integration + hope for best.

$$0 \leq x \leq 1$$

$$\sqrt{x} \leq y \leq 1$$

TYPE I



AS TYPE II REGION:

$$0 \leq y \leq 1$$

$$0 \leq x \leq y^2$$

$$\text{So } y=1 \quad x=y^2$$

$$I = \int_{y=0}^{y=1} \int_{x=0}^{x=y^2} \sqrt{y^3 + 1} dx dy$$


$$= \int_{y=0}^{y=1} \left[\sqrt{y^3 + 1} \right]_{x=0}^{x=y^2}$$

$$= \int_{y=0}^{y=1} \sqrt{y^3 + 1} y^2 dy$$

$$\begin{aligned} \text{SUB } u &= y^3 + 1 \\ du &= 3y^2 dy \end{aligned}$$

$$\begin{aligned} &= \frac{1}{3} \int_{u=1}^{u=2} \sqrt{u} du = \frac{1}{3} \left[\frac{2}{3} u^{3/2} \right]_{u=1}^{u=2} \\ &= \frac{2}{9} (2^{3/2} - 1) \end{aligned}$$

- (6) [10 pts] Let E be the solid in the first octant that is bounded by the planes $x+z=2$ and $2y+z=2$. Calculate $\iiint_E z \, dV$. Hint: It may be helpful to sketch E .




THESE 2 PLANE
meet in a line.
The points
 $(0, 0, 2)$ and
 $(2, 1, 0)$ lie
on the line

USE STICKS GO LEFT-RIGHT:

SHADOW OF STICKS ON xz -PLANE IS

$$0 \leq x \leq 2 \quad \left. \begin{array}{l} \\ \end{array} \right\} D$$

$$0 \leq z \leq 2-x$$



$$0 \leq y \leq 1 - \frac{z}{2} \quad \text{as } 2y+z=2 \text{ is RIGHT SURFACE}$$

$$\begin{aligned} \iiint_E z \, dV &= \int_0^2 \int_{x=0}^{x=2} \int_{z=0}^{z=2-x} z \left(1 - \frac{z}{2}\right) dz \, dx \\ &= \int_0^2 \int_{z=0}^{z=2-x} z \left(1 - \frac{z}{2}\right) dz \, dx = \int_{x=0}^{x=2} \left[\frac{z^2}{2} - \frac{z^3}{6} \right]_{z=0}^{z=2-x} \end{aligned}$$

$$= \int_0^2 \left(\frac{(2-x)^2}{2} - \frac{(2-x)^3}{6} \right) dx = \left[-\frac{(2-x)^3}{6} + \frac{(2-x)^4}{24} \right]_0^2 = \boxed{213}$$

(7) [10 pts]

- (a) Find $\iiint_E \sqrt{x^2 + y^2} dV$ where E is the solid above the xy -plane, below the paraboloid $z = 9 - x^2 - y^2$ and outside the cylinder $x^2 + y^2 = 4$.

Paraboloid and $z=0$ meet
in circle $x^2 + y^2 = 9$

CYL COORDS:

$$2 < r < \sqrt{3}$$

$$0 \leq \theta \leq 2\pi$$

$$0 \leq z \leq 9 - r^2$$

$$\iiint_E \sqrt{x^2 + y^2} dV = \int_0^{2\pi} \int_{r=2}^{\sqrt{3}} \int_{z=0}^{9-r^2} r^2 \cdot r dz dr d\theta$$

$$= 2\pi \int_{r=2}^{\sqrt{3}} r^3 (9 - r^2) dr = 2\pi \left[\frac{9}{3} r^3 - \frac{1}{5} r^5 \right]_2^{\sqrt{3}} = \boxed{\frac{3}{4} - \frac{3}{5} + \frac{32}{5}}$$

(b) Use a triple integral in spherical coordinates to find the volume of the sphere of radius R .

$$0 \leq \theta \leq 2\pi$$



$$0 \leq \phi \leq \pi$$

$$0 \leq \rho \leq R$$

$$dV = \rho^2 \sin \phi d\rho d\phi d\theta$$

$$VOL = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi} \int_{\rho=0}^R \rho^2 \sin \phi d\rho d\phi d\theta$$

$$= 2\pi \left(\int_0^{\pi} \sin \phi d\phi \right) \left(\int_0^R \rho^2 d\rho \right) = 2\pi \cdot (2) \left(\frac{\rho^3}{3} \right) \Big|_{\rho=0}^{R=1}$$




$$\begin{aligned} &= 2\pi \left[R^2 - \frac{35}{5} - 24 + \frac{32}{5} \right] \\ &= \frac{148\pi}{5} \end{aligned}$$

$$\boxed{\frac{4\pi R^3}{3}}$$


✓

!!

(8) [10 pts] Use the change of variables theorem to evaluate $\iint_R (x+y)^2 e^{x-y} dx dy$ where R is the parallelogram bounded by $x+y=2$, $x+y=5$, $x-y=-2$ and $x-y=1$.



$$\text{Let } u = x+y \\ v = x-y$$



$$u+v = 2x$$

$$\text{So } x = \frac{u+v}{2}$$

$$\frac{\partial(x,y)}{\partial(u,v)} = \text{DET} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix} = -\frac{1}{2}$$

$$u-v = 2y$$

$$\text{So } y = \frac{u-v}{2}$$

$$\left[\text{So } \iint_R (x+y)^2 e^{x-y} dx dy \right] = \iint_{u=2}^5 \int_{v=-2}^1 u^2 e^v \left| -\frac{1}{2} \right| dv du$$


$$= \frac{1}{2} \left(\int_2^5 u^2 du \right) \left(\int_{-2}^1 e^v dv \right)$$

$$= \frac{1}{2} \left(\frac{53}{3} - \frac{25}{3} \right) \left(e^1 - e^{-2} \right)$$

$$= \frac{117}{6} (e - e^{-2})$$

(9) [10 pts]

- (a) Let C be the semicircle $x^2 + y^2 = 9$ with $y \geq 0$ oriented counter-clockwise and let \mathbf{F} be the vector field $\mathbf{F} = -yi + xj$. Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$.



$$\vec{r}(t) = (3 \cos t, 3 \sin t), \quad 0 \leq t \leq \pi$$

$$\vec{r}'(t) = (-3 \sin t, 3 \cos t)$$

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^\pi \mathbf{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt \\ &= \int_0^\pi (-3 \sin t, 3 \cos t) \cdot (-3 \sin t, 3 \cos t) dt \\ &= \int_0^\pi 9 \sin^2 t + 9 \cos^2 t dt = \boxed{9\pi} \end{aligned}$$

- (b) Let C be the curve $y = \sqrt[3]{x+1}$ from $(-1, 0)$ to $(0, 1)$. Evaluate

$$\int_C P dx + Q dy$$

$$\frac{\partial Q}{\partial x} = 2x, \quad \frac{\partial P}{\partial y} = 2x$$

So $\frac{\partial Q}{\partial x} = \frac{\partial P}{\partial y}$ so $\mathbf{F} = \nabla f$ is conservative.

[Find f]

$$\frac{\partial f}{\partial x} = P \Rightarrow f(x, y) = \int (2xy + 3) dx = x^2y + 3x + g(y)$$


$$\frac{\partial f}{\partial y} = Q \Rightarrow f(x, y) = \int (x^2 + 10y) dy = x^2y + 5y^2 + h(x)$$

$$\text{So } f(x, y) = x^2y + 3x + 5y^2 + C \text{ works.}$$

$$\text{Pf: } \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(0, 1) - f(-1, 0) = 5 - (-3) = \boxed{8}$$

(10) [10 pts] Let D be the triangular domain with vertices $(0,0)$, $(0,4)$ and $(2,4)$. Let C be the boundary of D , oriented counter clockwise. Evaluate

$$I = \int_C (\sqrt{x^3 - 1} + 2xy^2)dx + (x^2y - e^y)dy.$$



D is Type I
 $0 \leq x \leq 2$
 $2x \leq y \leq 4$

By Green's Thm

$$\oint_C P dx + Q dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

$$\text{So } I = \iint_D (2xy - 4xy) dA = -2 \iint_D xy dA$$

$$= -2 \int_{x=0}^{x=2} \int_{y=2x}^{y=4} xy dy dx = -2 \int_{x=0}^{x=2} x \left[\frac{y^2}{2} \right]_{y=2x}^{y=4} dx$$

$$= -2 \int_{x=0}^{x=2} x (8 - 2x^2) dx \quad u = 8 - 2x^2 \\ du = -4x dx$$

$$= \frac{1}{2} \int_{u=8}^{u=0} u du = \left[\frac{u^2}{4} \right]_8^0 = \boxed{-16}$$